

To be published in the proceedings of “X-Ray Astronomy 2000”, Palermo, Italy

X-RAY BURSTERS A1742-294 AND SLX1744-299/300 NEAR THE GALACTIC CENTER

A.A.Lutovinov^{1,3,*}, S.A.Grebenev¹, M.N.Pavlinsky¹, R.A.Sunyaev^{1,2}

¹ Space Research Institute, Profsoyuznaya 84/32, 117810 Moscow, Russia

² Max-Planck-Institut für Astrophysik, Karl-Schwarzschild-Str. 1, 85740 Garching bei München, Germany

³ visiting INTEGRAL Science Data Center, ch. d’Ecogia 16, 1290 Vesoix, Switzerland

* e-mail: aal@hea.iki.rssi.ru

We present a study of two X-ray bursters in the Galactic Center field with the ART-P telescope onboard GRANAT. During 1990-1992 approximately 100 X-ray bursts were detected from the Galactic Center field and most of them were identified with the two persistent sources, A1742-294 and SLX1744-299/300. The burst recurrence times were found to be ~ 2.4 and ~ 4.6 hours, respectively, which is significantly shorter than was previously thought. We calculated the mean and maximum fluxes during the bursts and investigated their time profiles. We found that the shape of A1742-294 bursts correlates with the burst peak flux. The spectral analysis was carried out for the strongest burst, detected from A1742-294 on Oct 18, 1990.

1 INTRODUCTION

The first detailed X-ray map of the GC field was obtained by *EINSTEIN* in the soft energy band (≤ 4.5 keV). In the harder band (4-30 keV) this region was intensively studied only with the ART-P telescope, because other missions (*Spacelab-2*, *Spartan-1* and telescope TTM on the *MIR-KVANT* module) could not perform long observations for technical reasons. The total ART-P/*GRANAT* exposure time of the GC field observations is ~ 830 ks. Such a long exposure allowed to obtain the detailed X-ray map of the GC region, to investigate in detail emission from persistent sources, and to discover 4 new sources (Sunyaev et al., 1991, Pavlinsky et al., 1992).

This paper presents preliminary results of observations of two X-ray bursters A1742-294 and SLX1744-299/300 located $\sim 1^\circ$ from the GC. Both objects are regular sources of type I bursts. The time profiles of this type bursts depends on the energy band and are characterized by shorter decay time at high energies. Type I bursts are thought to result from thermonuclear flares on the surface of accreting neutron stars.

2 OBSERVATIONS

ART-P is a coded-mask telescope which is able to detect photons in the 3-60 keV energy band with the time resolution of 3.9 ms. The geometric area of the position-sensitive detector is 625 cm²; half of the detector is covered with opaque mask elements. The telescope field of view is 3°4×3°6, the energy resolution is ~25% at 5.9 keV and the instrument dead time is about 580 μ s. There are four identical telescope units on *GRANAT*. The telescope was described in detail by Sunyaev et al.(1990).

The observations were carried out in the “photon-by-photon” mode, in which coordinates, energy and arrival time of each photon were recorded into the buffer memory. The data are transferred from buffer to the main satellite memory when the buffer is filled. The data transfer happens each 150-200 sec and takes ~30 sec. This results in gaps between individual exposures in ART-P data sets.

3 RESULTS

The telescope ART-P detected 100 x-ray bursts in the GC field during 1990-1992 (Grebenev et al., 2000). Lewin et al.(1976) found that most part of X-ray bursts observed by *SAS-3* from the GC field originated from three sources – A1742-289, A1743-28 and A1742-294. During ART-P observations no bursts were detected from A1742-289 and A1743-28.

A1742-294 is the brightest persistent X-ray source in the GC field in the standard X-ray energy band. It is responsible for ~ 1/3 of the total X-ray emission from this region. The source persistent flux in the 3-20 keV energy band varied in the range 20–50 mCrab during ART-P observations. This flux corresponds to a luminosity of $(0.4 - 1) \times 10^{37}$ erg s⁻¹ assuming a distance of ~8.5 kpc. In total, 26 type I X-ray bursts with a typical duration of ~ (15 – 20) sec were identified with A1742-294 (see Table 1, fluxes are shown in the 3-20 keV energy band). In several observations, two bursts from this source were detected. It allowed to measure the burst recurrence time t_r ~2.4 hr. This value is several times smaller than that obtained by Lewin et al.(1976).

The other GC field source for which ART-P detected X-ray bursts was SLX1744-299/300. This source was discovered by *Spacelab-2* (Skinner et al., 1987) and *Spartan-1* (Kawai et al., 1988). In fact it is a double X-ray source, with components SLX1744-299 and SLX1744-300 separated by ~3', of which the latter is a burster. The angular resolution of ART-P, ~5', does not allow to separate fluxes of these two sources. The average flux from the double source measured by ART-P was equal to ~20 mCrab, which corresponds to the luminosity 4.4×10^{36} erg s⁻¹. During the ART-P observations 6 type I X-ray bursts were detected from this region (see Table 2, fluxes are shown in the 3-20 keV energy band). One of these bursts was extremely strong. The coordinates of this burst ($R.A.(1950) = 17^h44^m14.^s7$ and $Decl.(1950) = -29^\circ58'19''$) coincide with the SLX1744-299 position, while SLX1744-300 was outside the position error cycle (Pavlinsky et al., 1994). The SLX1744-299/300 bursts recurrence time was equal to ~4.6 hr.

The energy spectra of the persistent emission of both sources, A1742-294 and SLX1744-299/300, are very similar. They can be described by a thermal bremsstrahlung model with parameters in the range $kT_{br} \simeq (6.4 - 10.5)$ keV for A1742-294 and $kT_{br} \simeq (4.7 - 9.2)$ keV for SLX1744-299/300. The hydrogen column density N_H was fixed at a value of 6×10^{22} cm⁻². Typical persistent spectra of these sources measured on Sept 9, 1990 are shown in Fig.1. The

energy spectra of A1742-294 bursts can be modeled by the blackbody spectrum with temperatures in the range 1.4–2.7 keV; the mean value of the temperature is $kT_{\text{bb}} = 1.81 \pm 0.39$ keV. The mean black-body temperature for SLX1744-299/300 bursts was slightly higher, $kT_{\text{bb}} = 2.40 \pm 0.32$ keV.

The time profiles of A1742-294 depend on the source flux during the burst. The shape of the burst profile is almost “triangle”, i.e. the rise time is close to the decay time when the mean flux in the 3-20 keV energy was $\sim 0.4 \text{ phot} \cdot \text{cm}^{-2} \text{s}^{-1}$. When the burst flux in the same energy band was higher, $\sim 0.9 \text{ phot} \cdot \text{cm}^{-2} \text{s}^{-1}$, the burst profiles have a “classical” shape, i.e. a rapid rise of intensity followed by a smooth decay. To study this behavior in more detail, we constructed average profiles of weak and strong bursts. We rejected bursts which occurred in the beginning or the end of exposures. The bursts were normalized to the same peak intensity and aligned by maxima of their intensity. In Fig.2, we show the average profiles of weak and strong bursts in three energy bands. It is evident that the shapes of weak and strong bursts in the hard energy band (8-20 keV) are similar, while in the soft energy band (3-8 keV), the rise time of weak burst is longer than that for strong bursts.

The average profile of 4 bursts from SLX1744-299/300 is presented in different energy bands in Fig.3.

We carried out a spectral and timing analysis of the strongest X-ray burst detected on Oct 18, 1990 from A1742-294. The burst peak flux was ~ 1.5 Crab in the 3-20 keV energy band. The time profiles of this burst in four energy bands are shown in Fig.4 (the zero point corresponds to the burst maximum in the broad energy band 3-20 keV). It is clear that the burst maximum at high energies is reached later than at low energies, i.e. a rise time in the 12-16 keV energy band (~ 5 sec) is longer than that in the 4-8 keV energy band (~ 1 -2 sec). On the contrary, the exponential decay time decreases from 8.7 sec in the 4-8 keV energy band to 3.9 sec in the 8-12 keV band, 2.3 sec in the 12-16 keV band, and 1.5 sec in the 16-20 keV. No significant burst emission was detected in the harder energy band (≥ 20 keV). The evolution of the source X-ray luminosity in the 3-20 keV energy band and its best-fit blackbody temperature are shown in Fig.5. It is interesting to note that the highest temperature during the burst was reached at the moment of maximum count rate at high energies.

This work was supported by RBRF grants 98-02-17056, 99-02-17178 and 00-15-99297.

4 REFERENCES

- Grebenев S., Lutovinov A., Pavlinsky M. et al. // 2000, in press.
Kawai N. et al. // 1988, *Astroph. J.*, **330**, 130.
Lewin W., Hoffman J., Doty J. et al. // 1976, *MNRAS*, **177**, 83.
Pavlinsky M., Grebenев S., Sunyaev R. // 1992, *Sov. Astron. Lett.*, **18**, 217.
Pavlinsky M., Grebenев S., Sunyaev R. // 1994, *Astroph. J.*, **425**, 110.
Skinner G. et al. // 1987, *Nature*, **330**, 554.
Sunyaev R. et al. // 1990, *Adv. Space Res.*, **10**, 223.
Syunyaev, R., Pavlinskii, M., Churazov, E. et al. // 1991, *Sov. Astr. Lett.*, **17**, 42.

Table 1. X-ray bursts detected from A1742-294 by GRANAT/ART-P in 1990-1992.

Date	T_0 , UT	Flux, $\text{phot}\cdot\text{cm}^{-2}\text{s}^{-1}$	Flux max, $\text{phot}\cdot\text{cm}^{-2}\text{s}^{-1}$	Duration, s	Signifi- cance	Head
20.03.90	18 ^h 05 ^m 00 ^s	0.379±0.052	0.902±0.092	26	17.5	1
	20 01 29	0.311±0.044	0.962±0.095	18	16.4	1
24.03.90	18 32 42	0.572±0.068	1.197±0.109	24	16.6	1
	21 42 00	0.461±0.059	0.822±0.086	17	12.0	1
8.04.90	14 06 38	0.421±0.053	0.757±0.080	24	18.8	1
9.09.90	14 07 48	0.363±0.041	0.835±0.071	18	11.6	4
	15 53 21	0.386±0.043	0.804±0.069	17	16.0	4
29.09.90	13 32 38	0.164±0.018	0.279±0.026	30	8.5	4
5.10.90	15 31 44	0.208±0.030	0.294±0.039	32	12.0	4
	19 43 33	0.187±0.029	0.296±0.040	21	11.9	4
6.10.90	22 10 36	0.551±0.057	0.864±0.074	22	12.0	4
9.10.90	17 20 52	0.334±0.038	0.652±0.060	11	13.3	4
	19 06 11	0.373±0.042	0.831±0.070	15	17.8	4
10.10.90	14 35 48	0.369±0.041	0.608±0.058	12	15.3	4
	16 06 25	0.589±0.057	0.798±0.069	9	12.5	4
18.10.90	9 50 31	0.982±0.078	2.299±0.125	17	69.9	4
22.02.91	14 07 59	0.987±0.119	2.055±0.189	13	33.6	3
23.02.91	22 24 36	0.969±0.119	2.381±0.207	16	42.6	3
26.02.91	11 00 20	0.911±0.114	2.026±0.188	15	19.5	3
1.04.91	13 39 44	0.507±0.077	1.090±0.131	28	16.4	3
	15 28 57	0.705±0.099	1.413±0.155	19	18.5	3
8.04.91	13 16 25	0.654±0.091	1.601±0.164	22	13.0	3
15.10.91	19 59 08	0.626±0.095	1.710±0.178	14	17.9	3
18.10.91	11 44 04	0.592±0.086	1.321±0.147	15	24.6	3
21.02.92	12 52 53	0.755±0.103	2.453±0.213	21	39.9	3
2.03.92	1 43 22	0.929±0.120	2.322±0.207	15	29.8	3

Table 2. X-ray bursts detected from SLX1744-299/300 by GRANAT/ART-P in 1990-1992.

Date	Source	T_0 , UT	Flux, $\text{phot}\cdot\text{cm}^{-2}\text{s}^{-1}$	Flux max, $\text{phot}\cdot\text{cm}^{-2}\text{s}^{-1}$	Duration, s	Signifi- cance	Head
9.09.90	SLX1744-299/300	15 42 10	0.339±0.039	0.627±0.058	19	12.2	4
	SLX1744-299/300	20 05 21	0.529±0.053	0.940±0.076	16	13.8	4
10.09.90	SLX1744-299/300	00 52 25	0.270±0.033	0.657±0.061	15	17.2	4
9.10.90	SLX1744-299	19 19 58	0.720±0.064	2.742±0.138	170	86.3	4
18.10.90	SLX1744-299/300	11 07 49	0.583±0.056	1.211±0.088	21	17.1	4
23.02.91	SLX1744-299/300	21 35 50	0.391±0.063	0.930±0.119	15	9.1	3




FIG 1. Spectra of the persistent emission of A1742-294 (a) and SLX1744-299/300(b) obtained by GRANAT/ART-P on Sept 8, 1990. The best-fit models are shown by histograms. The solid lines represent the corresponding photon spectra.




FIG 2. The average profiles of weak (*left panel*) and strong (*right panel*) X-ray bursts detected from A1742-294.




FIG.3. The average SLX1744-299/300 X-ray burst profiles in different energy bands.




FIG.4. Time profiles of the strong X-ray burst detected by ART-P from A1742-294 on Oct 18, 1990.




FIG.5. Evolution of the A1742-294 luminosity in 3-20 keV energy band during the Oct 18, 1990 burst (*upper panel*). The persistent luminosity level is shown by dotted line. The evolution of corresponding best-fit black-body temperature is shown in the *bottom panel*.